




ACAAWS



African-Caribbean & African American Women's Study



Research supported by the Abuse Status and Health Effects among African Caribbean and African American Women (ACAAWS) study (J. Campbell & D. Campbell, Co-PI's) from the Caribbean Exploratory NCMHD Research Center of Excellence (CERC), University of the Virgin Islands, Grant # 5P20MD002286, National Institutes of Health, G. Callwood PI

Slide 1



DC1

Doris Campbell, 10/21/2010

Suicidality in African-Caribbean and African American Women Experiencing Intimate Partner Abuse

Doris Williams Campbell, PhD, ARNP, FAAN¹, Jacquelyn C. Campbell, PhD, RN, FAAN²
Desiree R. Bertrand, MSN, RN¹, Gloria B. Callwood, PhD, RN, FAAN¹,
Phyllis W. Sharps, PhD, RN, FAAN², Hossein Yarandi, PhD³, Richelle Bolyard, MHS²,
Catherine Coverston, PhD, RNC⁴, Kaitlan Gibbons, PsyD ©⁵,
Members of the ACAAWS Team


¹University of the Virgin Islands, ²Johns Hopkins University, ³Wayne State University,
⁴Brigham Young University, ⁵Argosy University





Background and Significance

- Suicide- Self killing
- Suicidality-No clear consensus on definition.
- One operational definition of contingent suicidality was: either the patient made suicide threats or statements that were linked to the admission decision
(Lambert, M.T (2002))
- Term suicidality not as clinically useful as more specific terminology (ideation, behavior, attempts, and suicide) Meyer et. al (2010)





Background and Significance

- Prior research links IPA and suicidality in low income African American women (Kaslow, Leiner, Reviere, Jackson, et al, 2002; Leiner, Compton, Houry & Kaslow, 2008)
- Limited studies of suicidality in Caribbean women
- Few studies examined risk factors associated with suicidality in underresourced communities, or culture and religious beliefs that may discourage suicide and promote self preservation among abused Caribbean women (McFarlane, Malacha, Gist, Watsdson, Batten, Hall and Smith, 2005; Kaslow, Thompson, Okun, Price, Young, Bender, Wyckoff, Goldin, Parker, 2001).




PURPOSE

- Describe the prevalence and mediators of suicidality in African Caribbean and African American women experiencing Intimate Partner Abuse (IPA) in the USVI and Baltimore City.
- Describe the association of suicidality with depression and post traumatic stress syndrome in African Caribbean and African American women experiencing IPV in the USVI and Baltimore.
- Describe resource and service utilization among African Caribbean and African women who experience IPV
- Describe pre-emptive interventions that might help African American and African Caribbean women cope effectively with suicidality.




Methods

- ❖ Comparative case (Abused) control (Non abused) study
- ❖ Eligibility criteria
 - ❖ Women aged 18-55 years
 - ❖ Self-identify as African Caribbean or African American
 - ❖ Report of a male sex partner in the past two years
- ❖ Women recruited from primary care, prenatal or family planning clinics
- ❖ Questionnaire administered on a touch screen computer with optional headphones - an important methodological strategy
 - ❖ For women of low literacy
 - ❖ For sensitive information
 - ❖ Alerts interviewer if high score on DA or suicidality
 - ❖ Allows increased recruitment among Spanish speaking
 - ❖ Recording in Spanish & English by USVI residents for increased cultural appropriateness
- ❖ Study period 12/1/09 to 09/2011



Case Status

- ❖ Intimate Partner Violence (IPV) in lifetime (case status) assessed using the Abuse Assessment Screen (McFarlane & Helton - www.nnvawi.org)
 - ❖ Pushed, slapped, hit, kicked, or physically hurt **&/OR**
 - ❖ Forced sex
 - Any of the above by current or former husband, boyfriend, or male or female partner
- ❖ **&/or** above cutoff on WEB (Women's Experiences of Battering – Hall-Smith)
 - ❖ Controlled, in fear of current or former husband, boyfriend, or male or female partner





MEASURES

- Suicidal thoughts
- Suicide attempts
- Self harm (self cutting)
- Mental & Physical QOL – from SF8
- Depression – CESD
- PTSD – brief PTSD screen

RECRUITMENT SITES

Baltimore City



RECRUITMENT SITES


Virgin Islands

St. Thomas



St. Croix





- RESULTS




Demographic Characteristics

	Baltimore City	US Virgin Islands	Overall
Age (In Years)	30.09 (SD=10.09)	29.28 (SD=8.55)	29.49 (DS=9.11)
Education			
Less than HS	28.18%	16.27%	20.38%
HS/GED	39.55%	39.71%	39.66%
Some College	17.73%	28.47%	24.76%
College Grad	14.55%	15.55%	15.20%
Marital Status			
Single	57.40%	43.33%	48.21%
Partnered	22.87%	34.29%	30.33%
Married	12.11%	16.67%	15.09%
Div./Wid./Other	7.62%	5.71%	6.38%
Employment	39.46%	52.14%	47.74%
Employed	60.54%	47.86%	52.26%
Unemployed			


PREVALENCE OF INTIMATE PARTNER VIOLENCE

	Baltimore City	US Virgin Islands	Overall
Type of IPV			
Lifetime Emotional Abuse	5.41%	7.89%	7.20%
Lifetime Physical/Sexual Abuse	32.48%	33.17%	32.98%
IPV in the Past 2 Years	21.14%	21.87%	21.66%
Lifetime Physical Abuse	13.45%	15.62%	15.04%
During Pregnancy			
Current Physical Abuse	6.60%	10.47%	9.09%
During Pregnancy			
Forced Sex in Past 2 Years	7.05%	9.63%	8.88%
Forced Sex in Past 2 Years By Intimate Partner	5.70%	6.88%	6.54%


Depression: Abused vs. Non-abused Women*




PTSD: Abused vs. Non-abused Women*




Co-morbid PTSD & Depression: Abused vs. Non-abused Women*





Depression & PTSD among Baltimore Cases




Depression and PTSD among USVI Cases





■ Baltimore ■ USVI





Abused Women

$c = 1.83$ ($p < 0.0001$), $OR = 6.25$

Suicide Thoughts


***Abused Women, relative to Non-Abused,
had 6.25 times higher odds having suicide thoughts.***

Abused Women


$c = 1.77$ ($p < 0.0001$), $OR = 5.88$

Suicide Attempts


***Abused Women, relative to Non-Abused,
had 5.88 times higher odds suicide attempts.***




Sobel Test: $Z = 6.63$, $P < 0.0001$, the mediator depression significantly carries the relationship of abused women and their suicide thoughts.



Sobel Test: $Z = 7.63$, $P < 0.0001$, the mediator depression significantly carries the relationship of abused women and their suicide attempts.



Sobel Test: $Z = 9.44$, $P < 0.0001$, the mediator PTSD significantly carries the relationship of abused women and their suicide thoughts.




Sobel Test: $Z = 8.92$, $P < 0.0001$, the mediator PTSD significantly carries the relationship of abused women and their suicide attempts.




SUICIDALITY: ODDS RATIO

- When age, income, education, employment, marital status, being pregnant, and children under 18 years of age were controlled for, relative to control group, women who had experienced abuse had
 - 2.12 times higher odds of engaging in self-cutting
 - 6.43 times higher odds of having suicide thoughts
 - 6.18 times higher odds of attempting suicide


RESOURCE UTILIZATION





CONCLUSIONS AND RECOMMENDATIONS

- At the IPV-Depression/PTSD link, depression mediated by many ways of coping, social support and spiritual support
- Ability to access resources is critical
- Routine universal screening for IPV in all health care settings recognizing that woman may present with many symptoms/complaints seemingly unrelated to IPV
- ACAAWS study protocol to assess for suicidality.



CONCLUSIONS AND RECOMMENDATIONS

- Assess for level of family functioning and social support. High levels of family functioning and support can mediate association between social environment and suicidality
- Clinicians need to explore among women victims, the presence of self-inflicted bodily self-harm
- Knowing that some coping styles or behaviors are protective against suicidality, practitioners should target these behaviors as risk reducing.



ACWS Team Members




United States Mainland

Jacquelyn Campbell, PhD, RN, FAAN
Phyllis Sharps PhD, RN, FAAN
Richelle Bolyard, MHS
Jamila Stockman, PhD, MPH
Marguerite Baty, PhD, MSN, MPH, RN
Akosoa McFadgion, MS, MSW, PhD student
Kaitlan Gibbons, PsyD (c)
Mary Paterno, MSN, CNM, RN, PhD student
Sharon O'Brien PhD
Sachi Mana-ay, BSN student
Jessica Draughon, MSN, RN, PhD student
Callie Simkoff, BSN
Gyasi Moscou-Jackson, MHS, BSN, RN
Chris Kunselman
Ayanna Johnson, BA
Ashley Chappell, BSN, RN
Lucine Francis, BSN
Naa Ayele Amponsah, BA
Hosseini Yarandi, PhD

US Virgin Islands

Doris Campbell, PhD, ARNP, FAAN
Gloria Callwood, PhD, RN, FAAN
Desiree Bertrand, MSN, RN
Charmayne M. Dunlop-Thomas, MS
Lorna Sutton, MPA
Tyra DeCastro
Alexandria Bradley, RN
Sally Browne, RN
Edris Evans, RN
Yvonne Francis, RN
Naomi Joseph, BSN student
Jennifer King, RN
Suzette Lettsume, MSPHN, RN
Julie Matthew, RN
Kenice Pemberton, ASN student
J'Nique Smith, BSN student
Jaslene Williams, MSW

- 
- THANK YOU!
 - THANK YOU!
 - QUESTIONS????
 - QUESTIONS????